

FLUBENDAZOLE 1%



Medicated Premixture

Domestic Poultry Wormer

Composition:

Each gram contains: 10 mg flubendazole

60 g tub (600 mg flubendazole)

This Medicated Premixture is produced from Flubenvet 5% w/w Premix for Medicated Feeding Stuff containing flubendazole 50mg/g, Vm 43877/4008, Marketing Authorisation Holder:

Kernfarm B.V.

De Corridor 14D

3621 ZB Breukelen, The Netherlands

Active against:

Flubendazole 1% Medicated Premixture is effective against gapeworm, large roundworm, caecal worm, hairworm, and gizzard worm in chickens, turkeys, and geese. Activity includes adult worms, larvae and eggs.

Dosage and administration

Dosage Instructions:

Medicated feed should be fed for 7 consecutive days. Incorporate the required amount of the product into the daily feed ration.

The daily ration should be fed as a "complete ration" during the 7 days. At the inclusion rates in the table below, for chicken and geese the ration will contain 30 ppm flubendazole and for turkeys 20 ppm.

Chicken and Geese	Turkeys
60 g of product (the whole tub) medicates 20 kg of feed	60 g of product (the whole tub) medicates 30 kg of feed
30 g of product medicates 10 kg of feed	18 g of product medicates 9 kg of feed
6 g of product medicates 2 kg of feed	6 g of product medicates 3 kg of feed

A level measure of the scoop included = 6 g of wormer. Use the scoop to measure the correct amount of product to mix into the feed as per the table above.

The dosage is dependent on the amount of feed that each bird consumes over the 7-day treatment period.

Larger birds eat more feed and receive more medication according to their weight.

Mixing advice:

The volume of product to be mixed into the feed is small. Mix very well for even distribution.

Initially, mix the required volume into a cup of feed. Mix very well. Add this to a bowl of feed and mix very well. Repeat twice more, every time into a larger volume of food. Mix well.

Treatment frequency:

Birds on the ground with known worm infestations are susceptible to re-infestation. In such cases, re-treatment with another 7-day course, after 3 weeks is recommended.

Flubendazole 1% Medicated Premixture has no adverse effect on egg laying hatchability.

Contra-indications, warnings, and precautions***Withdrawal periods:***

Birds must not be slaughtered for human consumption during treatment. Treated birds may be slaughtered for human consumption only after 7 days from the last treatment: Chickens, turkeys, geese: Meat: 7 days. There is no withholding period for chicken producing eggs for human consumption when the product is used at the recommended inclusion rates; Chickens eggs: zero days.

To be supplied only on veterinary prescription – from your veterinarian or a suitably qualified person.

Keep out of the reach and sight of children.

For Animal Treatment Only.

Store in tightly closed original container below 25 °C.

Operator warnings:

Accidental ingestion by humans should be avoided.

May cause sensitisation by skin contact.

May cause skin and eye irritation.

Avoid direct skin contact.

Wear overalls, safety glasses and impervious gloves when mixing and handling the product.

Wash affected parts if skin contact occurs.

If accidental eye contact occurs, immediately rinse thoroughly with water.

If the operations involve potential exposure to dust, wear either a disposable filter and half-mask respirator conforming to European Standard EN149, or a non-disposable respirator to European standard EN140 fitted with a filter to EN143.

Disposal Advice:

Dispose of empty containers in the domestic refuse. Used containers should not be recycled

Pack sizes:

60 g tub with 6 g scoop included

**Kernfarm**

LIVESTOCK PHARMA